HÆMATOPUS. Will. p. 297. *Bellon. Lib. III. p.* 203.

The Oyſter Catcher

WEighs one pound and two ounces; the Bill long, ſtreight and of a bright red colour, contracted near the baſis, and towards the end compreſs'd: the Irides of their Eyes are yellow, encompaſſed with a red Circle: the whole Head and Neck black, having a ſpot of white under the Eyes; all the under part of the Body, duſky white: the larger Quill-feathers are duſky black: the Tail is ſhort, black towards the End, and towards the Rump white: the upper part of the Body and Wings, brown, except a broad white line, which runs along the middle of each Wing; the Legs are long and thick, and of a reddiſh colour. It has only three fore-toes, wanting the back-toe. Their Feet are remarkably arm'd with a very rough ſcaly ſkin. In Rivers and Creeks near the Sea there are great quantities of Oyſter-banks, which at low Water are left bare: On theſe Banks of Oyſters do theſe Birds principally, if not altogether, ſubſiſt; Nature having not only formed their Bills ſuitable to the Work, but armed the Feet and Legs for a defence againſt the ſharp edges of the Oyſters. The Hens differ from the Cocks in not having the red circle round their Eyes, and their Bellies are of a more dirty white than in the Cocks: In the Maw of one was found nothing but indigeſted Oyſters.

This Bird ſeems to be the *Hæmatopus* of *Bellonius*, *Will. p.* 297. notwithſtanding there is ſome ſmall difference in their deſcription. I have ſeen them on the Sea Coaſts both of *Carolina* and the *Bahama* Iſlands.

Le Preneur d'Huiſtres.

CET oiſeau peſe une livre & deux onces. Son bec eſt long, droit & d'un rouge éclatant: il eſt ſerré vers ſa baſe, & applati par le bout. L'iris de ſes yeux eſt jaune, entouré d'un cercle rouge; toute ſa tête & ſon col ſont noirs: il a une tache blanche ſous les yeux; tout le deſſous de ſon corps eſt d'un blanc ſale: les grandes plumes de l'aile ſont d'un noir obſcur; la queue eſt courte, noire à l'extremité & blanche vers le croupion; le deſſus de ſon corps & de ſes ailes eſt brun, excepté une large raye blanche qui traverſe chaque aile par le milieu. Ses jambes ſont longues, épaiſſes & d'une couleur rougeatre. Il n'a que trois doits à chaque pied ſans ergot par derriere. Ses pieds ſont remarquables par la peau rude & écailleuſe dont ils ſont couverts. Dans les rivieres & les criques proche de la mer il y a grand nombre de bancs couverts d'Huitres, qui ſont à ſec lorſque la mer eſt baſſe; c'eſt ſur ces bancs que ces oiſeaux trouvent principalement leur ſubſiſtance, la nature leur ayant non ſeulement donné un bec formé de maniere à venir à bout de cet ouvrage, mais ayant auſſi orné leurs jambes & leurs pieds contre les bords tranchans des écailles d'Huitres. Les femelles different des mâles en ce qu'elles n'ont pas le cercle rouge autour des yeux, & que leur ventre eſt d'un blanc plus ſale que celui des mâles: on ne trouva dans le jabot d'un de ces oiſeaux que des Huitres qui n'étoient pas encore digerées.

Il ſemble que cet oiſeau eſt l'Hæmatopus de Bellonius, Will. p. 297. *quoiqu'il y ait quelque petite difference dans la deſcription de l'un & de l'autre. J'en ai vû ſur les côtes de la Caroline & des Iſles Bahama.*

Frutex Bahamenſis foliis oblongis ſucculentis, fructu ſubrotundo unicum nucleum continente.

THIS grows to the ſize of a ſmall Tree; the leaves ſtand by pairs on foot-ſtalks about an inch long; they are long, thick and ſucculent: at the ends of the ſtalks grow in pairs, and ſometimes ſingly, round flat Seed-veſſels, about the breadth of a ſhilling: the fruit is of the ſubſtance of a Bean, and, like that, divides in the middle: it is cover'd with a thin membrane of a pale green colour. I had no opportunity of ſeeing the Bloſſoms, tho' I was told they were very ſmall and white. The Bark of this Tree is uſed for tanning of Sole-leather.

IL croit juſques à la hauteur d'un petit arbre; ſes feuilles ſont attachées deux à deux par des pedicules d'environ un pouce de long: elles ſont longues, épaiſſes & ſucculentes. Au bout des tiges naiſſent par paires, ou ſeules, des capſules qui renferment les ſemences: elles ſont rondes, plattes, environ une pouce en diametre: le fruit reſſemble par ſa ſubſtance à une feve, &, comme elle, ſe diviſe en deux; il eſt couvert d'une membrane mince d'un vert pâle. Je n'ai pas eû la commodité de voir les fleurs; cependant on m'a dit qu'elles étoient blanches & très petites. L'écorce de cet arbre ſert à tanner le cuir dont on fait des ſemelles de ſoulier.